

Asymptotic Expansions of the Lauricella Hypergeometric Function F_D

Chelo Ferreira¹ and José L. López²

¹ *Departamento de Matemática Aplicada, Facultad de Ciencias,
Universidad de Zaragoza, 50013-Zaragoza, Spain. e-mail: cferrei@posta.unizar.es.*

² *Departamento de Matemática e Informática,
Universidad Pública de Navarra, 31006-Pamplona, Spain. e-mail: jl.lopez@unavarra.es.*

ABSTRACT

The Lauricella hypergeometric function $F_D^r(a, b_1, \dots, b_r; c; x_1, \dots, x_r)$ with $r \in \mathbb{N}$, is considered for large values of one variable: x_1 , or two variables: x_1 and x_2 . An integral representation of this function is obtained in the form of a generalized Stieltjes transform. Distributional approach is applied to this integral to derive four asymptotic expansions of this function in increasing powers of the asymptotic variable(s) $1 - x_1$ or $1 - x_1$ and $1 - x_2$. For certain values of the parameters a , b_i and c , two of these expansions also involve logarithmic terms in the asymptotic variable(s). For large x_1 , coefficients of these expansions are given in terms of the Lauricella hypergeometric function $F_D^{r-1}(a, b_2, \dots, b_r; c; x_2, \dots, x_r)$ and its derivative with respect to the parameter a , whereas for large x_1 and x_2 those coefficients are given in terms of $F_D^{r-2}(a, b_3, \dots, b_r; c; x_3, \dots, x_r)$ and its derivative. All the expansions are accompanied by error bounds for the remainder at any order of the approximation. Numerical experiments show that these bounds are considerably accurate.

2000 Mathematics Subject Classification: 41A60 33C65

Keywords & Phrases: Lauricella hypergeometric function, asymptotic expansions, distributional approach, generalized Stieltjes transforms.

1. Introduction

The Lauricella hypergeometric function $F_D^r(a, b_1, \dots, b_r; c; x_1, \dots, x_r)$ is defined by means of the multiple power series [7]

$$F_D^r(a, b_1, \dots, b_r; c; x_1, \dots, x_r) \equiv \sum_{m_1=0}^{\infty} \dots \sum_{m_r=0}^{\infty} \frac{(a)_{m_1+\dots+m_r} (b_1)_{m_1} \dots (b_r)_{m_r}}{(c)_{m_1+\dots+m_r} m_1! \dots m_r!} x_1^{m_1} \dots x_r^{m_r}, \quad |x_i| < 1, \quad i = 1, 2, \dots, r.$$

Two important particular cases of this function are the case $r = 1$: the Gauss hypergeometric function, and the case $r = 2$: the Appell function.

The Lauricella function has several applications in mathematical physics due to its role in the theory of Lie group representations [13] and its astrophysical and quantum chemical applications [21]. There is an extensive mathematical literature devoted to the study of these functions: Srivastava derived polynomial expansions of the Lauricella functions using expansions involving Bessel functions [22]. Integral formulas involving the Lauricella functions are established in [18]. The behavior of F_D^3 near certain of its logarithmic singularities is analyzed in [17]. Several formulas involving series and integrals of the Lauricella functions and their application to derive bilateral and bilinear generating functions of Jacobi, Laguerre and Sylvester polynomials are derived in [12]. In [1], a family of generalized elliptic-type integrals is introduced, which admits an explicit representation in terms of the Lauricella function. Carlson explains how the Lauricella hypergeometric function is connected with symmetric elliptic integrals and their advantages for numerical and symbolic integration [3]. The Lauricella functions have also probabilistic interpretations, which arise from the evaluation of certain product moments of some multivariate distributions. It is shown that those product moments may be expressed as transformations on the Lauricella functions [15]. Other specific probabilistic and statistic problems are studied in [11]. Generalized hypergeometric functions of several variables are needed to study Goursat-Darboux problems for third-order differential equations [4]. A comprehensive reference to the Lauricella function is [19].

Approximations of the Lauricella functions have been investigated by many authors: Complete power series expansions at $x_i = 0$, $i = 1, 2, 3, \dots, r-1$ of F_D^r in terms of products of the Gauss hypergeometric function and F_D^{r-1} have been obtained by Srivastava and Goyal [20], and later extended by Joshi [6]. Approximations by using branched continued fractions are studied in [14]. Zamel, Tuan and Kalla have obtained asymptotic expansions of F_D when one of its variables approaches 1 in terms of multiple series involving elementary functions [1]. On the other hand, asymptotic expansions of these functions for large values of the variables have not been exhaustively investigated. Complete convergent expansions of F_D^r have been obtained by Carlson using Mellin transforms techniques [2]. Although these expansions have an attractively simple structure, explicit computation of the terms of the expansions is not straightforward and the

upper bound on the truncation error is not quite satisfactory [[2], sec. 5]. Asymptotic approximations of a class of generalized hypergeometric functions, which includes F_D^r may be found in [4]. In this paper we consider the problem of finding complete asymptotic expansions of $F_D^r(a, b_1, \dots, b_r; c; x_1, \dots, x_r)$ for large values of one or two of its variables x_i . We attempt to obtain easy algorithms to compute the coefficients of these expansions as well as error bounds at any order of the approximation.

The starting point is the integral representation [[7], eq. (25)]

$$F_D^r(a, b_1, \dots, b_r; c; x_1, \dots, x_r) = \frac{\Gamma(c)}{\Gamma(c-a)\Gamma(a)} \int_0^1 \frac{s^{a-1}(1-s)^{c-a-1}ds}{(1-sx_1)^{b_1} \dots (1-sx_r)^{b_r}} \quad (1)$$

where $\Re(a) > 0$, $\Re(c-a) > 0$, $x_i \notin [1, \infty)$ if $b_i \geq 1$ for $i = 1, 2, 3, \dots, r$. This integral defines the analytical continuation of $F_D^r(a, b_1, \dots, b_r; c; x_1, \dots, x_r)$ to the cut complex x_i -planes $\mathcal{C} \setminus [1, \infty)$ [[23], p. 30, theorem 2.3].

The first step in the analysis is to write the aforementioned integral as a generalized Stieltjes transform. For that purpose, we perform the change of variable $s = (1+t)^{-1}$ in (1), obtaining:

$$F_D^r(a, b_1, \dots, b_r; c; x_1, \dots, x_r) = \frac{\Gamma(c)}{\Gamma(c-a)\Gamma(a)} \int_0^\infty \frac{f_r(t)}{(t+1-x_1)^{b_1}} dt, \quad (2)$$

or

$$F_D^r(a, b_1, \dots, b_r; c; x_1, \dots, x_r) \equiv \frac{\Gamma(c)}{\Gamma(c-a)\Gamma(a)} \int_0^\infty \frac{f_{r-1}(t)}{(t+1-x_1)^{b_1}(t+1-x_2)^{b_2}} dt, \quad (3)$$

where

$$f_{r-1}(t) \equiv \frac{t^{c-a-1}(1+t)^{b_1+b_2+\dots+b_r-c}}{\prod_{k=3}^r (t+1-x_k)^{b_k}}, \quad f_r(t) \equiv \frac{f_{r-1}(t)}{(t+1-x_2)^{b_2}}. \quad (4)$$

Then, up to a factor, the Lauricella function F_D^r is a generalized Stieltjes transform of $f_{r-1}(t)$ or $f_r(t)$. For $\Re(c-a) > 0$, the function $f_{r-1}(t)$ is a locally integrable function on $[0, \infty)$ and satisfies

$$f_{r-1}(t) = \sum_{k=0}^{n-1} \frac{A_k}{t^{k-b_1-b_2+a+1}} + f_n^{r-1}(t), \quad (5)$$

where

$$A_k \equiv \sum_{m_1=0}^k \sum_{m_2=0}^{m_1} \dots \sum_{m_{r-2}=0}^{m_{r-3}} \binom{\sum_{j=1}^r b_j - c}{k - m_1} \binom{-b_3}{m_1 - m_2} \dots \binom{-b_{r-1}}{m_{r-1} - m_{r-2}} \times \binom{-b_r}{m_{r-2}} (1-x_r)^{m_{r-2}} \prod_{i=1}^{r-3} (1-x_{i+2})^{m_i - m_{i+1}} \quad (6)$$

and $f_n^{r-1}(t) = \mathcal{O}(t^{-n+b_1+b_2-a-1})$ when $t \rightarrow \infty$.

On the other hand, for $\Re(c-a) > 0$ and $x_2 - 1 \notin \mathbb{R}^+ \cup \{0\}$ if $\Re b_2 \geq 1$, $f_r(t)$ is a locally integrable function on $[0, \infty)$ and satisfies

$$f_r(t) = \sum_{k=0}^{n-1} \frac{B_k}{t^{k-b_1+a+1}} + f_n^r(t), \quad (7)$$

where

$$B_k \equiv \sum_{j=0}^k A_{k-j} \binom{-b_2}{j} (1-x_2)^j \quad (8)$$

and $f_n^r(t) = \mathcal{O}(t^{-n+b_1-a-1})$ when $t \rightarrow \infty$.

Classical asymptotic methods do not apply to integrals (2) and (3). But the asymptotic methods based on the distributional approach [24] [[25], chap. 6], [8], [9] may be generalized in order to be applied to these integrals. This generalization is performed in [5], [10], and also in this paper. In section 2, we summarise the main theorems obtained in [5] and [10] and derive new theorems that we will need in this paper. In section 3 we apply those theorems to obtain asymptotic expansions of integrals (2) and (3) with error bounds, for both: large x_1 and fixed x_j , $j \neq 1$ and large x_1 and x_2 and fixed x_j , $j \neq 1, 2$. Several numerical examples are shown as illustrations.

2. Asymptotic expansions of generalized Stieltjes transforms

Let $f(t)$ be a locally integrable function on $[0, \infty)$ which satisfies

$$f(t) = \sum_{k=K}^{n-1} \frac{a_k}{t^{k+s}} + f_n(t), \quad (9)$$

where $K \in \mathbb{Z}$, $0 < \Re s \leq 1$, $\{a_k, k = K, K+1, K+2, \dots\}$ is a sequence of complex numbers and $f_n(t) = \mathcal{O}(t^{-n-s})$ when $t \rightarrow \infty$.

Then, asymptotic expansions (including error bounds) of the generalized Stieltjes transforms of $f(t)$,

$$S_f(w; z) \equiv \int_0^\infty \frac{f(t)}{(t+z)^w} dt, \quad S_f(w_1, w_2; z) \equiv \int_0^\infty \frac{f(t)}{(t+xz)^{w_1}(t+yz)^{w_2}} dt, \quad (10)$$

for large z and fixed x and y are given in [[25], chap. 6], [8] and [9] for real w , w_1 , w_2 and s and complex x , y , z . In [5] and [10], these theorems are generalized to the case of complex w , w_1 , w_2 and s . Therefore, in the following pages, we consider that the parameters w , w_1 , w_2 , x , y and z are complex and that $f(t)$ is a locally integrable function on $[0, \infty)$ which satisfies (9). In the following pages, we use the notation introduced in [25].

2.1. Asymptotic expansion of $S_f(w; z)$ and $S_f(w_1, w_2; z)$ for large z

Asymptotic expansions of $S_f(w; z)$ for large z are given in the following two theorems proved in [[10], theorems 1,2 and 3].

Theorem 1. *Let $f(t)$ be a locally integrable function on $[0, \infty)$ which satisfies (9) with $0 < \Re s \leq 1$, $s \neq 1$. Then, for $z \in \mathbb{C} \setminus \mathbb{R}^- \cup \{0\}$, $\Re(s + w) + K > 1$ and $n = 1, 2, 3, \dots$,*

$$\begin{aligned} \int_0^\infty \frac{f(t)}{(t+z)^w} dt &= \sum_{k=K}^{n-1} \frac{(-1)^k \pi a_k \Gamma(w+s+k-1)}{\Gamma(s+k) \Gamma(w) \sin(\pi s) z^{w+s+k-1}} + \\ &\quad \sum_{k=0}^{n-1} \frac{(-1)^k (w)_k M[f; k+1]}{k! z^{k+w}} + R_n(w; z), \end{aligned} \quad (11)$$

where empty sums must be understood as zero and $(w)_k$ denotes the Pochhammer symbol. In this formula, $M[f; z]$ denotes the Mellin transform of f : $\int_0^\infty t^{z-1} f(t) dt$ or its analytic continuation. The remainder term is defined by

$$R_n(w; z) \equiv (w)_n \int_0^\infty \frac{f_{n,n}(t) dt}{(t+z)^{n+w}}, \quad (12)$$

where $f_{n,n}(t)$ is defined recursively by

$$f_{n,0}(t) \equiv f_n(t), \quad f_{n,k+1}(t) \equiv - \int_t^\infty f_{n,k}(u) du = \frac{(-1)^{k+1}}{k!} \int_t^\infty (u-t)^k f_n(u) du. \quad (13)$$

The remainder term verifies $R_n(w; z) = \mathcal{O}(z^{-n-w-s+1})$ when $z \rightarrow \infty$.

Theorem 2. *Let $f(t)$ be a locally integrable function on $[0, \infty)$ which satisfies (9) with $s = 1$. Then, for $z \in \mathbb{C} \setminus \mathbb{R}^- \cup \{0\}$, $\Re w + K > 0$ and $n = 1, 2, 3, \dots$,*

$$\begin{aligned} \int_0^\infty \frac{f(t)}{(t+z)^w} dt &= \sum_{k=K}^{-1} a_k \frac{\Gamma(w+k) \Gamma(-k)}{\Gamma(w) z^{w+k}} + \sum_{k=0}^{n-1} \frac{(-1)^k (w)_k}{k! z^{k+w}} \times \\ &\quad [a_k (\log(z) - \gamma - \psi(k+w)) + b_{k+1}] + R_n(w; z), \end{aligned} \quad (14)$$

where empty sums must be understood as zero and for $k = 0, 1, 2, \dots$, the coefficients b_{k+1} are given by

$$b_{k+1} \equiv \lim_{z \rightarrow k} \left[M[f; z+1] + \frac{a_k}{z-k} \right] + a_k (\gamma + \psi(k+1)), \quad (15)$$

The remainder term $R_n(w; z)$ is given in (12) and verifies $R_n(w; z) = \mathcal{O}(z^{-n-w} \log z)$ when $z \rightarrow \infty$.

To obtain now asymptotic expansions of $S_f(w_1, w_2; z)$ for large z we use the same techniques that we used in [10]. We denote by $\mathcal{S}$ the space of rapidly decreasing functions and by $\langle \Lambda, \varphi \rangle$ the image of a tempered distribution Λ acting over a function $\varphi \in \mathcal{S}$.

Since $f(t)$ in (9)-(10) is a locally integrable function on $[0, \infty)$, it defines a distribution $\mathbf{f}$:

$$\langle \mathbf{f}, \varphi \rangle \equiv \int_0^\infty f(t) \varphi(t) dt.$$

The distributions associated with t^{-k-s} , $k = 0, 1, 2, \dots, n-1$ are given by [[25], chap. 5], [10]:

$$\begin{aligned} \langle \mathbf{t}^{-\mathbf{k}-s}, \varphi \rangle &\equiv \frac{1}{(s)_k} \int_0^\infty t^{-s} \varphi^{(k)}(t) dt & \text{if } 0 < \Re s < 1, \\ \langle \mathbf{t}^{-\mathbf{k}-s}, \varphi \rangle &\equiv \frac{1}{(i\Im s)_{k+1}} \int_0^\infty t^{-i\Im s} \varphi^{(k+1)}(t) dt & \text{if } 1 \neq s = 1 + i\Im s, \end{aligned}$$

where $(s)_k$ denotes the Pochhammer symbol of s , and

$$\langle \mathbf{t}^{-\mathbf{k}-1}, \varphi \rangle \equiv -\frac{1}{k!} \int_0^\infty \log(t) \varphi^{(k+1)}(t) dt.$$

To assign a distribution to the function $f_n(t)$ introduced in (9), we use the recursive definition of $f_{n,n}(t)$ given in (13) [[25], chap. 5], [10]:

$$\langle \mathbf{f}_n, \varphi \rangle \equiv (-1)^n \langle \mathbf{f}_{n,n}, \varphi^{(n)} \rangle \equiv (-1)^n \int_0^\infty f_{n,n}(t) \varphi^{(n)}(t) dt.$$

Following the same steps in [10], we define a particular function of $\mathcal{S}$:

$$\varphi_\eta(t) \equiv \frac{e^{-\eta t}}{(t+xz)^{w_1}(t+yz)^{w_2}} \in \mathcal{S},$$

where $xz \notin \mathbb{R}^- \cup \{0\}$ if $\Re w_1 \geq 1$, $yz \notin \mathbb{R}^- \cup \{0\}$ if $\Re w_2 \geq 1$ and $\eta > 0$. In the following lemma we calculate the image of the tempered distributions defined above and of the delta distribution acting over $\varphi_\eta(t)$, and take the limit $\eta \rightarrow 0$.

Lemma 1. *Let $f(t)$ verify (9). Then, for $0 < \Re s \leq 1$, $k = 0, 1, 2, \dots$ and $n = 1, 2, 3, \dots$, the following identities hold,*

$$\lim_{\eta \rightarrow 0} \langle \mathbf{f}, \varphi_\eta \rangle = \int_0^\infty \frac{f(t)}{(t+xz)^{w_1}(t+yz)^{w_2}} dt \quad \text{for } \Re(s+w_1+w_2)+K > 1.$$

$$\lim_{\eta \rightarrow 0} \langle \delta, \varphi_\eta^{(k)} \rangle = \frac{(-1)^k}{z^{k+w_1+w_2}} \sum_{j=0}^k \binom{k}{j} \frac{(w_1)_j (w_2)_{k-j}}{x^{w_1+j} y^{w_2+k-j}},$$

$$\begin{aligned} \lim_{\eta \rightarrow 0} \langle \mathbf{t}^{-s}, \varphi_\eta^{(k)} \rangle &= \frac{\Gamma(1-s)\Gamma(k+w_1+w_2+s-1)}{\Gamma(w_1+w_2)z^{k+w_1+w_2+s-1}} \frac{(-1)^k}{x^{w_1+k+s-1} y^{w_2}} \times \\ &F\left(\begin{matrix} 1-s-k, w_2 \\ w_1+w_2 \end{matrix} \middle| 1-\frac{x}{y}\right) \quad \text{for } \Re(s+w_1+w_2)+k > 1, \quad s \neq 1, \end{aligned}$$

where $F\left(\begin{smallmatrix}\alpha, \beta \\ \delta\end{smallmatrix} \middle| z\right) \equiv {}_2F_1(\alpha, \beta, \delta; z)$ denotes the Gauss hypergeometric function,

$$\begin{aligned} \lim_{\eta \rightarrow 0} \langle \mathbf{log}(\mathbf{t}), \varphi_{\eta}^{(k+1)} \rangle = & \frac{(-1)^{k+1}}{(k+w_1+w_2)z^{k+w_1+w_2}} \sum_{j=0}^{k+1} \binom{k+1}{j} \frac{(w_1)_j (w_2)_{k+1-j}}{x^{w_1+j-1} y^{w_2+k+1-j}} \times \\ & \left[(\log(xz) - \gamma - \psi(k+w_1+w_2)) F\left(\begin{smallmatrix}1, k+1+w_2-j \\ k+1+w_1+w_2\end{smallmatrix} \middle| 1 - \frac{x}{y}\right) + \right. \\ & \left. F'\left(\begin{smallmatrix}1, k+1+w_2-j \\ k+1+w_1+w_2\end{smallmatrix} \middle| 1 - \frac{x}{y}\right) \right], \quad \text{for } \Re(s+w_1+w_2) > 0, \end{aligned}$$

where $F'\left(\begin{smallmatrix}\alpha, \beta \\ \delta\end{smallmatrix} \middle| z\right) \equiv \frac{d}{d\alpha} F\left(\begin{smallmatrix}\alpha, \beta \\ \delta\end{smallmatrix} \middle| z\right)$ and

$$\lim_{\eta \rightarrow 0} \langle \mathbf{f}_{\mathbf{n}, \mathbf{n}}, \varphi_{\eta}^{(n)} \rangle = (-1)^n \sum_{j=0}^n \binom{n}{j} \int_0^{\infty} \frac{(w_1)_j (w_2)_{n-j} f_{n,n}(t)}{(t+xz)^{j+w_1} (t+yz)^{n-j+w_2}} dt$$

for $\Re(s+w_1+w_2)+n > 1$.

Proof. It is a straightforward generalization of the proof of [[5], lemma 4] from real to complex values of s , w_1 and w_2 . $\square$

Using the aforementioned lemma 1 and lemmas 1 and 2 of [10], we prove the following theorems.

Theorem 3. Let $f(t)$ be a locally integrable function on $[0, \infty)$ which satisfies (9) with $0 < \Re s \leq 1$, $s \neq 1$. Then, for $xz, yz \in \mathcal{C} \setminus \mathbb{R}^- \cup \{0\}$, $\Re w_1, \Re w_2 > 0$, $\Re(s+w_1+w_2)+K > 1$ and $n = 1, 2, 3, \dots$,

$$\int_0^{\infty} \frac{f(t)}{(t+xz)^{w_1} (t+yz)^{w_2}} dt = \sum_{k=K}^{n-1} \frac{A_k}{z^{k+w_1+w_2+s-1}} + \sum_{k=0}^{n-1} \frac{B_k}{z^{k+w_1+w_2}} + R_n(w_1, w_2; z), \quad (16)$$

where empty sums must be understood as zero. The coefficients A_k , B_k and C_k are defined by

$$A_k \equiv a_k \frac{\Gamma(1-s-k)\Gamma(w_1+w_2+s+k-1)}{\Gamma(w_1+w_2)x^{w_1+s+k-1}y^{w_2}} F\left(\begin{smallmatrix}1-s-k, w_2 \\ w_1+w_2\end{smallmatrix} \middle| 1 - \frac{x}{y}\right),$$

and

$$B_k \equiv \frac{(-1)^k M[f; k+1]}{k!} \sum_{j=0}^k \binom{k}{j} \frac{(w_1)_j (w_2)_{k-j}}{x^{w_1+j} y^{k+w_2-j}}.$$

The remainder term is defined by

$$R_n(w_1, w_2; z) \equiv \sum_{j=0}^n \binom{n}{j} (w_1)_j (w_2)_{n-j} \int_0^{\infty} \frac{f_{n,n}(t) dt}{(t+xz)^{j+w_1} (t+yz)^{n-j+w_2}}, \quad (17)$$

where $f_{n,n}(t)$ is defined in (13) and verifies $R_n(w_1, w_2; z) = \mathcal{O}(z^{-n-w_1-w_2-s+1})$ when $z \rightarrow \infty$.

Proof. The proof of formulas (16)-(17) is a straightforward generalization of the proof of theorem 3 in [5] from real to complex parameters. It uses previous lemma 1 instead of lemma 4 as used there, which is valid only for real parameters. The proof of the asymptotic character of expansion (16) is a straightforward generalization of the proof of [[5], theorem 5] to the case of complex parameters (theorem 5 in [5] is proved only for real parameters). $\square$.

Theorem 4. Let $f(t)$ be a locally integrable function on $[0, \infty)$ which satisfies (9) with $s = 1$. Then, for $xz, yz \in \mathcal{C} \setminus \mathbb{R}^- \cup \{0\}$, $\Re w_1, \Re w_2 > 0$, $\Re(w_1 + w_2) + K > 0$ and $n = 1, 2, 3, \dots$,

$$\int_0^\infty \frac{f(t)}{(t+xz)^{w_1}(t+yz)^{w_2}} dt = \sum_{k=K}^{-1} \frac{A_k}{z^{w_1+w_2+k}} + \sum_{k=0}^{n-1} \frac{(-1)^k}{k! z^{k+w_1+w_2}} \left[B_k (\log(xz) - \gamma - \psi(k+w_1+w_2)) + B'_k + C_k \right] + R_{n,s}(w_1, w_2; z), \quad (18)$$

where empty sums must be understood as zero,

$$A_k \equiv a_k \frac{\Gamma(-k)\Gamma(w_1+w_2+k)}{\Gamma(w_1+w_2)x^{w_1+k}y^{w_2}} F\left(\begin{matrix} -k, w_2 \\ w_1+w_2 \end{matrix} \middle| 1 - \frac{x}{y}\right),$$

$$B_k \equiv \frac{a_k}{(k+w_1+w_2)} \sum_{j=0}^{k+1} \binom{k+1}{j} \frac{(w_1)_j (w_2)_{k+1-j}}{x^{w_1+j-1} y^{w_2+k+1-j}} F\left(\begin{matrix} 1, k+1+w_2-j \\ k+1+w_1+w_2 \end{matrix} \middle| 1 - \frac{x}{y}\right),$$

$$B'_k \equiv \frac{a_k}{(k+w_1+w_2)} \sum_{j=0}^{k+1} \binom{k+1}{j} \frac{(w_1)_j (w_2)_{k+1-j}}{x^{w_1+j-1} y^{w_2+k+1-j}} F'\left(\begin{matrix} 1, k+1+w_2-j \\ k+1+w_1+w_2 \end{matrix} \middle| 1 - \frac{x}{y}\right),$$

and

$$C_k \equiv b_{k+1} \sum_{j=0}^k \binom{k}{j} \frac{(w_1)_j (w_2)_{k-j}}{x^{w_1+j} y^{k+w_2-j}},$$

where b_{k+1} is given in (15). The remainder term $R_{n,s}(w_1, w_2; z)$ is given in (17) and verifies $R_n(w_1, w_2; z) = \mathcal{O}(z^{-n-w_1-w_2} \log z)$ when $z \rightarrow \infty$.

Proof. The proof of the previous formulas is a straightforward generalization of the proof of theorem 4 in [5] from real to complex parameters. Again, it uses previous lemma 1 instead of lemma 4 as used there. And again, the proof of the asymptotic character of expansion (18) is a straightforward generalization of the proof of [[5], theorem 5] to the case of complex parameters. $\square$.

2.2. Error bounds

The following two propositions show that, if $\exists c_n > 0$ such that $|f_n(t)| \leq c_n t^{-n-\Re s} \forall t \in [0, \infty)$ then, bounds for the remainder terms in the expansions above can be calculated in terms of the constant c_n .

Proposition 1. *If, for $0 < \Re s < 1$, the remainder $f_n(t)$ in the expansion (9) of the function $f(t)$ satisfies the bound $|f_n(t)| \leq c_n t^{-n-\Re s} \forall t \in [0, \infty)$ for some positive constant c_n then, the remainder $R_n(w; z)$ in the expansion (11) satisfies*

$$|R_n(w; z)| \leq \frac{c_n \pi (|w|)_n \Gamma(n + \Re w + \Re s - 1) h(z, w)}{\Gamma(n + \Re s) \Gamma(n + \Re w) |\sin(\pi \Re s)| |z|^{n + \Re w + \Re s - 1}} \times F \left(\begin{matrix} 1 - \Re s, n + \Re s + \Re w - 1 \\ (n + \Re w + 1)/2 \end{matrix} \middle| \sin^2 \left(\frac{\text{Arg}(z)}{2} \right) \right) \quad (19)$$

and the remainder $R_{n,s}(w_1, w_2; z)$ in the expansion (16) satisfies

$$|R_n(w_1, w_2; z)| \leq \frac{c_n \pi (|w_1| + |w_2|)_n \Gamma(n + \Re(w_1 + w_2 + s) - 1) h(xz, w_1) h(yz, w_2)}{\Gamma(n + \Re s) \Gamma(n + \Re w_1 + \Re w_2) |\sin(\pi \Re s)| |vz|^{n + \Re(w_1 + w_2 + s) - 1}} \times F \left(\begin{matrix} 1 - \Re s, n + \Re(s + w_1 + w_2) - 1 \\ (n + \Re w_1 + \Re w_2 + 1)/2 \end{matrix} \middle| \frac{1}{2} \left(1 - \frac{u}{|vz|} \right) \right),$$

where

$$v \equiv \text{Min}\{|x|, |y|\}, \quad u \equiv \text{Min}\{\Re(xz), \Re(yz)\} \quad (20)$$

and

$$h(z, w) \equiv \begin{cases} 1 & \text{if } \text{Arg}(z) \Im w \geq 0 \\ e^{|\text{Arg}(z) \Im w|} & \text{if } \text{Arg}(z) \Im w < 0. \end{cases} \quad (21)$$

Proof. (19) is proved in [[10], proposition 1]. The second bound is obtained using (19), the inequalities $|t + xz|^2, |t + yz|^2 \geq t^2 + 2ut + |vz|^2$ in the definition (17) of $R_{n,s}(w_1, w_2; z)$, formula [[16], p. 309, eq. 7] and the equality

$$\sum_{k=0}^n \binom{n}{k} (|w_1|)_k (|w_2|)_{n-k} = (|w_1| + |w_2|)_n. \quad (22)$$

□

Proposition 2. *If, for $\Re s = 1$, each remainder $f_n(t)$ in the expansion (9) of the function $f(t)$ satisfies the bound $|f_n(t)| \leq c_n t^{-n-1} \forall t \in [0, \infty)$ for some positive constant c_n then, the remainder $R_n(w; z)$ in the expansions (11) and (14) satisfies*

$$|R_n(w; z)| \leq \frac{\bar{c}_n \pi (|w|)_n \Gamma(n + \Re w - 1/2) h(z, w)}{\Gamma(n + 1/2) \Gamma(n + \Re w) |z|^{n + \Re w - 1/2}} \times F \left(\begin{matrix} 1/2, n + \Re w - 1/2 \\ (n + \Re w + 1)/2 \end{matrix} \middle| \sin^2 \left(\frac{\text{Arg}(z)}{2} \right) \right) \equiv \mathbf{R}_n^{(1)}(w; z), \quad (23)$$

where $\bar{c}_n \equiv \text{Max}\{c_n, c_{n-1} + |a_{n-1}|\}$ and

$$|R_n(w; z)| \leq \frac{(|w|)_n}{|z|^{n+\Re w}} \left\{ \frac{\epsilon(c_{n-1} + |a_{n-1}|) + c_n}{(n-1)!\Theta(z, \epsilon)^{n+\Re w}} + \frac{c_n}{n!} \left| 1 + \frac{\epsilon}{z} \right|^{-n-\Re w} \left[\log |z| + \frac{(n + \Re w)[(2\epsilon + \Re z + |\Re z|)(|z|^{-1} - 1) + (|\Re z| - \Re z) \log |z|]}{2(n + \Re w + 1)|z + \epsilon|} H_1 + \frac{4\epsilon + \Re z + |\Re z| - 2\epsilon|z|}{2\epsilon(n + \Re w + 1)|z|} H_0 + \frac{2|\epsilon + z|H_{-1}}{\epsilon((n + \Re w)^2 - 1)|z|} \right] \right\} h(z, w) \equiv \mathbf{R}_n^{(2)}(w; z), \quad (24)$$

where ϵ is an arbitrary positive number,

$$H_m \equiv F \left(\begin{matrix} 2 - m, n + \Re w + m \\ (n + \Re w + 3)/2 \end{matrix} \middle| \sin^2 \left(\frac{\text{Arg}(z + \epsilon)}{2} \right) \right) \quad (25)$$

and

$$\Theta(z, \epsilon) \equiv \begin{cases} 1 & \text{if } \Re z \geq 0 \\ |\sin(\text{Arg}(z))| & \text{if } \epsilon \geq -\Re z > 0. \\ |1 + \epsilon/z| & \text{if } -\Re z > \epsilon > 0. \end{cases} \quad (26)$$

For large z and fixed n , the optimum value for ϵ is given approximately by

$$\epsilon^2 = \frac{c_n}{n(c_{n-1} + |a_{n-1}|)} \left[\frac{2H_{-1}}{(n + \Re w)^2 - 1} + \frac{(\Re z + |\Re z|)H_0}{2(n + \Re w + 1)|z|} \right]. \quad (27)$$

The remainder $R_n(w_1, w_2; z)$ in expansions (16) and (18) satisfies

$$|R_n(w_1, w_2; z)| \leq \mathbf{R}_n^{(i)}(w_1 + w_2; xz) + \mathbf{R}_n^{(i)}(w_1 + w_2; yz), \quad (28)$$

With either $i = 1$ or $i = 2$. If x, y, z, w, w_1 and w_2 are positive real numbers, then

$$|R_n(w; z)| \leq [n\epsilon(c_{n-1} + |a_{n-1}|) + c_n(S_n(z, \epsilon, w) + T_n(z, \epsilon, w))] \frac{(w)_n}{n!z^{n+w}}, \quad (29)$$

where ϵ is again an arbitrary positive number,

$$S_n(z, \epsilon, w) \equiv \text{Min} \left\{ \frac{nz [(\epsilon + z)^{n+w-1} - z^{n+w-1}]}{\epsilon(n + w - 1)(\epsilon + z)^{n+w-1}}, \psi(n + 1) + \gamma \right\}$$

and

$$T_n(z, \epsilon, w) \equiv \frac{z^{n+w}}{(n + w)(\epsilon + z)^{n+w}} F \left(n + w, 1; n + w + 1; \frac{z}{\epsilon + z} \right).$$

For large z and fixed n , the optimum value for ϵ is given by

$$\epsilon = \frac{c_n}{n(c_{n-1} + |a_{n-1}|)}. \quad (30)$$

The remainder $R_n(w_1, w_2; z)$ in expansions (16) and (18) satisfies the bound (29) replacing w by $w_1 + w_2$ and z by vz , with $v \equiv \text{Min}\{|x|, |y|\}$.

Proof. The bounds (23)-(27) for $R_n(w; z)$ are obtained in [[10], proposition 2].

On the other hand, the bound (28) is obtained using the inequality $|t + xz|^{-\Re w_1} |t + yz|^{-\Re w_2} \leq |t + xz|^{-\Re w_1 - \Re w_2} + |t + yz|^{-\Re w_1 - \Re w_2}$ in the definition (17) of $R_n(w_1, w_2; z)$ and formulas (22), (23) and (24).

The bounds for $R_n(w; z)$ and $R_n(w_1, w_2; z)$ for real positive x, y, w, w_1, w_2 and z have been obtained in [[9], propositions 2 and 4]. $\square$

In the following two propositions, two families of functions $f(t)$ are given, which verify the bound $|f_n(t)| \leq c_n t^{-n-\Re s}$ required in the aforementioned propositions 1 and 2. Moreover, the constant c_n is easily obtained from $f(t)$. These propositions are proved in lemmas 5 and 6 and corollaries 1 and 2 in [5].

Proposition 3. *If $f(t)$ verifies (9) with $\Re s > 0$ and the function $g(u) \equiv u^{-s-K} f(u^{-1})$ is a bounded analytic function in the region W of the complex z -plane comprised by the points situated at a distance $< \sigma$ from the positive real axis (see fig. 1), then $R_{n,s}(w; z)$ and $R_{n,s}(w_1, w_2; z)$ satisfy the bounds given in propositions 1 and 2 with $c_n = Cr^{-n}$, where C is a bound of $|g(z)|$ in W and $0 < r < \sigma$. Moreover, the expansions given in theorems 1 and 2 are convergent when the parameter $|z|$ is longer than the inverse of the width of the region W : when $\sigma|z| \geq 1$ if $\Re w < 1$ or $\sigma|z| > 1$ if $\Re w \geq 1$. The expansions given in theorems 3 and 4 are convergent when the parameter $|vz|$, with $v \equiv \text{Min}\{|x|, |y|\}$, is longer than the inverse of the width of that region: when $\sigma|vz| \geq 1$ if $\Re w_1 + \Re w_2 < 1$ or $\sigma|vz| > 1$ if $\Re w_1 + \Re w_2 \geq 1$.*

For $\Re s = 1$, the convergence of these expansions requires also that $\lim_{n \rightarrow \infty} n^{w-1} a_n z^{-n} = 0$ or $\lim_{n \rightarrow \infty} n^{w_1+w_2-1} a_n (vz)^{-n} = 0$ respectively.




Figure 1. Analyticity region W for the function $g(z)$ considered in proposition 3.

Proposition 4. *If the expansion (9) of $f(t)$ verifies the error test, then $R_n(w; z)$ and $R_{n,s}(w_1, w_2; z)$ satisfy the bounds given in propositions 1 and 2 replacing c_n by $|a_n|$ and c_{n-1} by 0. Moreover, the expansions given in theorems 1 and 2 are convergent when the coefficients a_n in the asymptotic expansion (9) verify $\lim_{n \rightarrow \infty} n^{w-1} a_n z^{-n} = 0$. The expansions given in theorems 3 and 4 are convergent when $\lim_{n \rightarrow \infty} n^{w_1+w_2-1} a_n (vz)^{-n} = 0$, $v \equiv \text{Min}\{|x|, |y|\}$.*

3. Asymptotic expansions of the Lauricella function F_D^r

In order to obtain asymptotic expansions of $F_D^r(a, b_1, \dots, b_r; c; x_1, \dots, x_r)$ for $|x_1| \rightarrow \infty$ and/or $|x_2| \rightarrow \infty$ we just apply theorems 1-4 to the integrals (2) or (3). Error bounds for the remainders are obtained from propositions 3 and 4, because the functions $f_{r-1}(t)$ and $f_r(t)$ which define F_D^r in (2) and (3) belong to the class of functions considered in those propositions.

Corollary 1. For $\Re a > 0$, $\Re(c-a) > 0$, $1+a-b_1 \notin \mathbb{Z}$, $|\text{Arg}(z)| < \pi$ and $x_i - 1 \notin \mathbb{R}^+ \cup \{0\}$ if $\Re b_i \geq 1$ for $2 \leq i \leq r$,

$$F_D^r(a, b_1, \dots, b_r; c; 1-z, x_2, \dots, x_r) = \frac{\Gamma(c)}{\Gamma(c-a)\Gamma(a)} \left\{ \sum_{k=0}^{n-1} \frac{\Gamma(b_1+k)}{\Gamma(b_1)} \frac{(-1)^k C_k}{k! z^{k+b_1}} + \frac{\pi}{\Gamma(b_1) \sin(\pi s)} \sum_{k=0}^{n-K-1} \frac{(-1)^k \Gamma(k+a)}{\Gamma(k-b_1+a+1)} \frac{B_k}{z^{k+a}} + R_n(a, b_1, \dots, b_r; c; z, x_2, \dots, x_r) \right\}, \quad (31)$$

where $K \equiv \text{Int}(\Re(1-b_1+a))$ and the coefficients B_k are defined in (8). Coefficients C_k are given by

$$C_k \equiv \frac{\Gamma(k+c-a)\Gamma(a-k-b_1)}{\Gamma(c-b_1) \prod_{j=2}^r (1-x_j)^{b_j}} \times F_D^{r-1} \left(k-a+c; b_2, \dots, b_r; c-b_1, \frac{x_2}{x_2-1}, \dots, \frac{x_r}{x_r-1} \right), \quad (32)$$

where $s \equiv \text{Fr}(\Re(1+a-b_1)) + i\Im(1+a-b_1)$.

If $\Re(1+a-b_1) \notin \mathbb{Z}$ and $n \in \mathbb{N}$, a bound for the remainder is given by

$$|R_n(a, b_1, \dots, b_r; c; z, x_2, \dots, x_r)| \leq \frac{c_n \pi (|b_1|)_n \Gamma(n + \Re(b_1 + s) - 1)}{\Gamma(n + \Re s) \Gamma(n + \Re b_1) |\sin(\pi \Re s)|} \times \quad (33)$$

$$\frac{h(z, b_1)}{|z|^{n+\Re(b_1+s)-1}} F \left(\begin{matrix} 1 - \Re s, n + \Re(s + b_1) - 1 \\ (n + \Re b_1 + 1)/2 \end{matrix} \middle| \sin^2 \left(\frac{\text{Arg}(z)}{2} \right) \right),$$

where $h(z, b_1)$ was defined in (21) and we can take $c_n = |B_{n-K}|$ if the following conditions over the parameters hold:

$$a, c, b_1, \dots, b_r \in \mathbb{R}, \quad \sum_{j=1}^r b_j \leq c, \quad b_j \geq 0, \quad \Re(1-x_j) > 0, \quad j = 2, 3, 4, \dots, r. \quad (34)$$

In any case, we can take $c_n = Cr^{-n}$, where

$$C \geq \sup_{u \in W} \left| (1+u)^{b_1+b_2+\dots+b_r-c} \left(\prod_{j=2}^r (1+(1-x_j)u)^{-b_j} \right) \right|, \quad (35)$$

W is the region considered in Proposition 3 for $g(u) = u^{b_1-a-1}f_{r-1}(u^{-1})$ with

$$0 < r < \text{Min} \{1, |1 - x_2|^{-1}\xi(b_2), \dots, |1 - x_r|^{-1}\xi(b_r)\}, \quad (36)$$

$$\xi(b) \equiv \begin{cases} 1 & \text{if } b \notin \mathbb{Z}^- \cup \{0\} \\ +\infty & \text{if } b \in \mathbb{Z}^- \cup \{0\}. \end{cases}$$

On the other hand, if $\Re(1 + a - b_1) \in \mathbb{Z}$ and $n \in \mathbb{N}$, two bounds for the remainder are given by

$$|R_n(a, b_1, \dots, b_r; c; z, x_2, \dots, x_r)| \leq \frac{\bar{c}_n \pi(|b_1|)_n \Gamma(n + \Re b_1 - 1/2) h(z, b_1)}{\Gamma(n + 1/2) \Gamma(n + \Re b_1) |z|^{n + \Re b_1 - 1/2}} \times \\ F \left(\begin{matrix} 1/2, n + \Re b_1 - 1/2 \\ (n + \Re b_1 + 1)/2 \end{matrix} \middle| \sin^2 \left(\frac{\text{Arg}(z)}{2} \right) \right) \equiv \mathbf{R}_n^{(1)}(B_n, b_1; z) \quad (37)$$

and

$$|R_n(a, b_1, \dots, b_r; c; z, x_2, \dots, x_r)| \leq \frac{(|b_1|)_n}{|z|^{n + \Re b_1}} \left\{ \frac{c_n}{n!} \left| 1 + \frac{\epsilon}{z} \right|^{-n - \Re b_1} \left[\log |z| + \right. \right. \\ \left. \frac{(n + \Re b_1)[(2\epsilon + \Re z + |\Re z|)(|z|^{-1} - 1) + (|\Re z| - \Re z) \log |z|]}{2(n + \Re b_1 + 1)|z + \epsilon|} H_1 + \right. \\ \left. \frac{4\epsilon + \Re z + |\Re z| - 2\epsilon|z|}{2\epsilon(n + \Re b_1 + 1)|z|} H_0 + \frac{2|\epsilon + z|}{\epsilon((n + \Re b_1)^2 - 1)|z|} H_{-1} \right] + \\ \left. \frac{\epsilon(c_{n-1} + |B_{n-K-1}|) + c_n}{(n-1)!\Theta(z, \epsilon)^{n + \Re b_1}} \right\} h(z, b_1) \equiv \mathbf{R}_n^{(2)}(Cr^{-n}, B_n, b_1; z). \quad (38)$$

In these formulas, $\bar{c}_n = \text{Max}\{|B_{n-K}|, |B_{n-K-1}|\}$ with $c_n = |B_{n-K}|$ and $c_{n-1} = 0$ if conditions (34) hold. In any case, we can take $\bar{c}_n = \text{Max}\{c_n, c_{n-1} + |B_{n-K-1}|\}$, with $c_n = Cr^{-n}$ given above. In (38), ϵ is an arbitrary positive number, $\Theta(z, \epsilon)$ is given in (26) and H_k is given in (25) setting $w = b_1$. For large z and fixed n , the optimum value for ϵ is given approximately by (27) setting $w = b_1$. Moreover, the expansion (31) is convergent when $\text{Max}\{|1 - x_2|\xi(b_2)^{-1}, \dots, |1 - x_r|\xi(b_r)^{-1}, 1\} < |z|$.

Proof. To obtain the expansion (31), we just apply theorem 1 to the integral (2) with $f(t) = f_r(t)$ given in (4), $a_k = B_{k-K}$ given in (8), $w = b_1$ and s and K given above. After the change of variable $t = u(1 - u)^{-1}$, the mellin transform of $f_r(t)$ reads

$$M[f_r; k + 1] = \left(\prod_{i=2}^r (1 - x_i)^{-b_i} \right) \int_0^1 \frac{u^{k+c-a-1}}{(1-u)^{k+b_1-a+1}} \prod_{i=2}^r \left(1 + \frac{x_i}{1-x_i} u \right)^{-b_i} du.$$

Then, the first term in (32) follows from (1).

If (34) holds, then, by [[8], lemmas 3 and 4], the function $f_r(t)$ verifies the error test. Therefore, by proposition 4, the remainder in the expansion (31) verifies the bounds given in propositions 1 and 2 with $c_n = |B_{n-K}|$, $c_{n-1} = 0$. In any case, by proposition

3, the remainder in the expansion (31) verifies the bounds given in propositions 1 and 2 with $c_n = Cr^{-n}$, C and r verifying (35) and (36) respectively. Therefore, the bounds (33), (37) and (38) hold.

Now introduce the bound

$$|B_n| \leq C \binom{\Re(b_1 - c)}{n} [\text{Max}\{1, |1 - x_2|\xi(b_2)^{-1}, \dots, |1 - x_r|\xi(b_r)^{-1}\}]^n,$$

where C is independent of n , in (33) and (37). Then, using (36), we obtain that $\lim_{n \rightarrow \infty} R_n(a, b_1, \dots, b_r; c; z, x_2, \dots, x_r) = 0$ if $\text{Max}\{|1 - x_2|\xi(b_2)^{-1}, \dots, |1 - x_r|\xi(b_r)^{-1}, 1\} < |z|$. $\square$

Corollary 2. For $\Re a, \Re(c - a) > 0$, $x_i - 1 \notin \mathbb{R}^+ \cup \{0\}$ if $\Re b_i \geq 1$ for $2 \leq i \leq r$, $1 + a - b_1 \in \mathbb{Z}$ and $|\text{Arg}(z)| < \pi$,

$$\begin{aligned} F_D^r(a, b_1, \dots, b_r; c; 1 - z, x_2, \dots, x_r) &= \frac{\Gamma(c)}{\Gamma(c - a)\Gamma(a)} \left\{ \sum_{k=0}^{n-1} \frac{(-1)^k (b_1)_k}{k! z^{k+b_1}} \times \right. \\ &\quad [B_{k+b_1-a}(\log(z) - \gamma - \psi(k + b_1)) + C_k] + \sum_{k=0}^{b_1-a-1} B_k \frac{\Gamma(k + a)\Gamma(b_1 - k - a)}{\Gamma(b_1)z^{k+a}} + \\ &\quad \left. R_n(a, b_1, \dots, b_r; c; z, x_2, \dots, x_r) \right\}, \end{aligned} \quad (39)$$

where the coefficients B_k are given in (8) and the coefficients C_k are given by

$$\begin{aligned} C_k &\equiv \frac{(-1)^{k+b_1-a-1}\Gamma(k + c - a)}{\Gamma(c - b_1)(k + b_1 - a)! \prod_{i=2}^r (1 - x_i)^{b_i}} \left\{ [\psi(k + c - a) - \right. \\ &\quad \left. \psi(k + b_1 - a + 1)] F_D^{r-1} \left(k - a + c, b_2, \dots, b_r; c - b_1; \frac{x_2}{x_2 - 1}, \dots, \frac{x_r}{x_r - 1} \right) + \right. \\ &\quad \left. F_D^{r-1'} \left(k - a + c, b_2, \dots, b_r; c - b_1; \frac{x_2}{x_2 - 1}, \dots, \frac{x_r}{x_r - 1} \right) \right\} + \\ &\quad B_{k+b_1-a}(\gamma + \psi(k + 1)). \end{aligned} \quad (40)$$

where $F_D^{r-1'}$ represents the derivative of the Lauricella function F_D^{r-1} with respect to the first parameter.

For $n \in \mathbb{N}$, two bounds for the remainder are given by (37) and (38) in corollary 1 replacing B_{n-K} by B_{n-a+b_1} . The expansion (39) is convergent if $\text{Max}\{|1 - x_2|\xi(b_2)^{-1}, \dots, |1 - x_r|\xi(b_r)^{-1}, 1\} < |z|$, where $\xi(b_i)$ is defined in (36).

Proof. To obtain the expansion (39), just apply theorem 2 to the integral (2) with $f(t) = f_r(t)$ given in (4), $s = 1$, $K = a - b_1$, $a_k = B_{k+b_1-a}$ and $w = b_1$.

On the other hand, the coefficients B_k in the expansion (7) of $f_r(t)$ may be written

$$B_k = \frac{1}{k!} \frac{d^k}{dt^k} \left[t^{b_1-a-1} f_r(t^{-1}) \right]_{t=0}.$$

Using the Cauchy formula for the derivative of an analytic function, we obtain

$$B_{k+b_1-a} = \frac{d^{k+b_1-a}}{dt^{k+b_1-a}} \left[\frac{t^k(1-t)^{b_1-a-1}}{(k+b_1-a)!} f_r \left(\frac{t}{1-t} \right) \right]_{t=1}. \quad (41)$$

The coefficient C_n in (39) is just b_{n+1} given by (15) with $a_n = B_{n+b_1-a}$. The Mellin transform $M[f_r; z+1]$ in formula (15) is given by

$$\frac{\Gamma(z+c-a)\Gamma(a-z-b_1)}{\Gamma(c-b_1)\prod_{j=2}^r(1-x_j)^{b_j}} F_D^{r-1} \left(z-a+c, b_2, \dots, b_r; c-b_1; \frac{x_2}{x_2-1}, \dots, \frac{x_r}{x_r-1} \right),$$

Then, when $z \rightarrow n$, there are two singular terms in the limit in (15): $B_{n+b_1-a}/(z-n)$ and $\Gamma(a-z-b_1)$. Setting $z = n + \eta$, expanding these terms at $\eta = 0$ and using (41) we obtain (40).

The bounds (37) and (38) are obtained as in corollary 1 (but using only proposition 2). $\square$

Corollary 3. For $\Re a > 0$, $\Re(c-a) > 0$, $1+a-b_1-b_2 \notin \mathbb{Z}$, $|\text{Arg}(xz)| < \pi$ and $|\text{Arg}(yz)| < \pi$,

$$\begin{aligned} F_D^r(a, b_1, \dots, b_r; c; 1-xz, 1-yz, x_3, \dots, x_r) &= \frac{\Gamma(c)}{\Gamma(c-a)\Gamma(a)} \left\{ \sum_{k=0}^{n-1} \frac{(-1)^k B_k}{k! z^{k+b_1+b_2}} + \right. \\ &\quad \sum_{k=0}^{n-K-1} \frac{\Gamma(b_1+b_2-a-k)\Gamma(a+k)}{\Gamma(b_1+b_2)x^{a+k-b_2}y^{b_2}z^{k+a}} A_k F \left(\begin{matrix} b_1+b_2-a-k, b_2 \\ b_1+b_2 \end{matrix} \middle| 1-\frac{x}{y} \right) + \\ &\quad \left. R_n(a, b_1, \dots, b_r; c; xz, yz, x_3, \dots, x_r) \right\}, \end{aligned} \quad (42)$$

where $K \equiv \text{Int}(\Re(1+a-b_1-b_2))$ and the coefficients A_k are defined in (6). Coefficients D_k are given by

$$\begin{aligned} B_k &\equiv \sum_{j=0}^k \binom{k}{j} \frac{(b_1)_j (b_2)_{k-j}}{x^{b+j} y^{k+c-j}} \frac{\Gamma(k+c-a)\Gamma(-k+a-b_1-b_2)}{\Gamma(c-b_1-b_2)\prod_{i=3}^r(1-x_i)^{b_i}} \times \\ &\quad F_D^{r-2} \left(k+c-a, b_3, \dots, b_r; c-b_1-b_2; \frac{x_3}{x_3-1}, \dots, \frac{x_r}{x_r-1} \right), \end{aligned} \quad (43)$$

where $s \equiv \text{Fr}(\Re(1-b_1-b_2+a)) + i\Im(1-b_1-b_2+a)$.

If $\Re(1+a-b_1-b_2) \notin \mathbb{Z}$ and $n \geq 0$, a bound for the remainder is given by

$$\begin{aligned} |R_n(a, b_1, \dots, b_r; c; xz, yz, x_3, \dots, x_r)| &\leq \\ &\frac{c_n \pi (|b_1| + |b_2|)_n \Gamma(n + \Re(b_1 + b_2 + s) - 1) h(xz, b_1) h(yz, b_2)}{\Gamma(n + \Re s) \Gamma(n + \Re(b_1 + b_2)) |\sin(\pi \Re s)| |vz|^{n + \Re(b_1 + b_2 + s) - 1}} \times \\ &F \left(\begin{matrix} 1 - \Re s, n + \Re(s + b_1 + b_2) - 1 \\ (n + \Re(b_1 + b_2) + 1)/2 \end{matrix} \middle| \frac{1}{2} \left(1 - \frac{u}{|vz|} \right) \right), \end{aligned} \quad (44)$$

where $h(z, w)$ was defined in (21) and u and v were defined in (20).

In formula (44) we can take $c_n = |A_{n-K}|$ if

$$a, c, b_1, \dots, b_r \in \mathbb{R}, \quad \sum_{j=1}^r b_j \leq c, \quad b_j \geq 0, \quad \Re(1 - x_j) > 0, \quad j = 3, 4, 5, \dots, r. \quad (45)$$

In any case, we can take $c_n = C$, where

$$C > \sup_{u \in W} \left| (1 + u)^{\sum_{j=1}^r b_j - c} \left(\prod_{j=3}^r (1 + (1 - x_j)u)^{-b_j} \right) \right| \quad (46)$$

and W is the region considered in proposition 3 for $g(u) = u^{b_1+b_2-a-1} f_{r-1}(u^{-1})$.

On the other hand, if $\Re(1 + a - b_1 - b_2) \in \mathbb{Z}$ and $n \geq 0$, the remainder in the expansion (42) satisfies

$$|R_n(a, b_1, \dots, b_r; c; xz, yz, x_3, \dots, x_r)| \leq \mathbf{R}_n^{(1)}(A_n, b_1 + b_2; 1 - xz) + \mathbf{R}_n^{(1)}(A_n, b_1 + b_2; 1 - yz) \quad (47)$$

or

$$|R_n(a, b_1, \dots, b_r; c; xz, yz, x_3, \dots, x_r)| \leq \mathbf{R}_n^{(2)}(C, A_n, b_1 + b_2; 1 - xz) + \mathbf{R}_n^{(2)}(C, A_n, b_1 + b_2; 1 - yz) \quad (48)$$

where $\mathbf{R}_n^{(1)}$ and $\mathbf{R}_n^{(2)}$ were defined in (37) and (38). Moreover, the expansion (42) is convergent if $v|z| > \max\{|1 - x_3|\xi(b_3)^{-1}, \dots, |1 - x_r|\xi(b_r)^{-1}, 1\}$.

Proof. To obtain the expansion (42), just apply theorem 3 to the integral (3) with $f(t) = f_{r-1}(t)$ given in (4), $a_k = A_{k-K}$ given in (6), $w_1 = b_1$, $w_2 = b_2$ and s and K given above. The calculation of coefficient C_k in formula (16) of theorem 3 requires the calculation of the Mellin transform $M[f_{r-1}; k+1]$. After trivial manipulations, and using (1) we obtain

$$M[f_{r-1}; k+1] = \frac{\Gamma(k+c-a)\Gamma(-k+a-b_1-b_2)}{\Gamma(c-b_1-b_2) \prod_{i=3}^r (1-x_i)^{b_i}} \times F_D^{r-2} \left(k+c-a, b_3, \dots, b_r; c-b_1-b_2; \frac{x_3}{x_3-1}, \dots, \frac{x_r}{x_r-1} \right)$$

and (43) follows.

If (45) holds, then, by [[8], lemmas 3 and 4], the function $f_{r-1}(t)$ verifies the error test. Therefore, by proposition 4, the remainder in the expansion (42) verifies the bounds given in propositions 1 and 2 with $c_n = |A_{n-K}|$, $c_{n-1} = 0$. In any case, by proposition 3, the remainder in the expansion (42) verifies the bounds given in formula (28) of proposition 2 with $w_1 = b_1$, $w_2 = b_2$, $r = 1$ and $c_n = C$, C verifying (46). Therefore, the bound (47) holds.

Finally, using the same argument that we used at the end of the proof of the corollary 1, we obtain that, if $\text{Min}\{|xz|, |yz|\} > \text{Max}\{|1 - x_3|\xi(b_3)^{-1}, \dots, |1 - x_r|\xi(b_r)^{-1}, 1\}$, then $\lim_{n \rightarrow \infty} R_n^{(i)} = 0$ and therefore, $\lim_{n \rightarrow \infty} R_n(a, b_1, \dots, b_r; c; xz, yz, x_3, \dots, x_r) = 0$. $\square$

Corollary 4. For $\Re a > 0$, $\Re(c - a) > 0$, $1 + a - b_1 - b_2 \in \mathbb{Z}$, $|\text{Arg}(xz)| < \pi$ and $|\text{Arg}(yz)| < \pi$,

$$\begin{aligned} F_D^r(a, b_1, \dots, b_r; c; 1 - xz, 1 - yz, x_3, \dots, x_r) &= \frac{\Gamma(c)}{\Gamma(c - a)\Gamma(a)} \left\{ \sum_{k=0}^{n-1} \frac{(-1)^k C_{k+1}}{k! z^{k+b_1+b_2}} + \right. \\ &\quad \sum_{k=0}^{b_1+b_2-a-1} \frac{\Gamma(-k + b_1 + b_2 - a)\Gamma(k + a)A_k}{\Gamma(b_1 + b_2)x^{k+a-b_2}y^{b_2}z^{a+k}} F\left(\begin{matrix} b_1 + b_2 - a - k, b_2 \\ b_1 + b_2 \end{matrix} \middle| 1 - \frac{x}{y}\right) + \\ &\quad \sum_{k=0}^{n-1} \frac{(-1)^k A_{k+b_1+b_2-a}}{k!(k + b_1 + b_2)z^{k+b_1+b_2}} [D_k(\log(xz) - \gamma - \psi(k + b_1 + b_2)) + D'_k] + \\ &\quad \left. R_n(a, b_1, \dots, b_r; c; xz, yz, x_3, \dots, x_r) \right\}, \end{aligned} \quad (49)$$

where coefficients A_k are given in (6) and coefficients D_k , D'_k and C_{k+1} are given by

$$\begin{aligned} D_k &\equiv \sum_{j=0}^{k+1} \binom{k+1}{j} \frac{(b_1)_j (b_2)_{k+1-j}}{x^{b_1+j-1} y^{b_2+k+1-j}} F\left(\begin{matrix} 1, k+1+b_2-j \\ k+1+b_1+b_2 \end{matrix} \middle| 1 - \frac{x}{y}\right), \\ D'_k &\equiv \sum_{j=0}^{k+1} \binom{k+1}{j} \frac{(b_1)_j (b_2)_{k+1-j}}{x^{b_1+j-1} y^{b_2+k+1-j}} F'\left(\begin{matrix} 1, k+1+b_2-j \\ k+1+b_1+b_2 \end{matrix} \middle| 1 - \frac{x}{y}\right) \end{aligned}$$

and

$$\begin{aligned} C_{k+1} &\equiv \left\{ \frac{\Gamma(k + c - a)(-1)^{k+b_1+b_2-a-1}}{(k + b_1 + b_2 - a)! \Gamma(c - b_1 - b_2) \prod_{i=3}^r (1 - x_i)^{b_i}} \times \right. \\ &\quad \left[(\psi(k + c - a) - \psi(k + b_1 + b_2 - a + 1)) \times \right. \\ &\quad \left. F_D^{r-2}\left(k + c - a, b_3, \dots, b_r; c - b_1 - b_2; \frac{x_3}{x_3 - 1}, \dots, \frac{x_r}{x_r - 1}\right) + \right. \\ &\quad \left. F_D^{r-2'}\left(k + c - a, b_3, \dots, b_r; c - b_1 - b_2; \frac{x_3}{x_3 - 1}, \dots, \frac{x_r}{x_r - 1}\right) \right] + \\ &\quad \left. A_{k+b_1+b_2-a}(\gamma + \psi(k + 1)) \right\} \sum_{j=0}^k \binom{k}{j} \frac{(b_1)_j (b_2)_{k-j}}{x^{b_1+j} y^{k+b_2-j}}. \end{aligned} \quad (50)$$

respectively.

The remainder in the expansion (49) satisfies the bounds (47) and (48). Moreover, the expansion (49) is convergent if $\text{Min}\{|xz|, |yz|\} > \text{Max}\{|1 - x_3|\xi(b_3)^{-1}, \dots, |1 - x_r|\xi(b_r)^{-1}, 1\}$.

Proof. To obtain the expansion (49), just apply theorem 4 to the integral (3) with $f(t) = f_{r-1}(t)$ given in (4), $s = 1$, $K = a - b_1 - b_2$, $a_k = A_{k+b_1+b_2-a}$, $w_1 = b_1$ and $w_2 = b_2$.

On the other hand, the coefficients A_k in the expansion (5) of $f_{r-1}(t)$ may be written

$$A_k = \frac{1}{k!} \frac{d^k}{dt^k} \left[t^{b_1+b_2-a-1} f_{r-1}(t^{-1}) \right]_{t=0}.$$

Using the Cauchy formula for the derivative of an analytic function, we obtain

$$A_{k+b+c-a} = \frac{d^{k+b_1+b_2-a}}{dt^{k+b_1+b_2-a}} \left[\frac{t^k (1-t)^{b_1+b_2-a-1}}{(k+b_1+b_2-a)!} f_{r-1} \left(\frac{t}{1-t} \right) \right]_{t=1}. \quad (51)$$

Coefficients C_{n+1} in (49) are given by b_{n+1} in (15) with $a_n = A_{n+b_1+b_2-a}$. The Mellin transform $M[f_{r-1}; z+1]$ in formula (15) is given by

$$\frac{\Gamma(z+c-a)\Gamma(a-z-b_1-b_2)}{\Gamma(c-b_1-b_2) \prod_{i=3}^r (1-x_i)^{b_i}} F_D^{r-2} \left(z+c-a, b_3, \dots, b_r; c-b_1-b_2; \frac{x_3}{x_3-1}, \dots, \frac{x_r}{x_r-1} \right).$$

Then, when $z \rightarrow n$, there are two singular terms in the limit in (15): $A_{n+b_1+b_2-a}/(z-n)$ and $\Gamma(a-z-b_1-b_2)$. Setting $z = n + \eta$, expanding these terms at $\eta = 0$ and using (51) we obtain (50).

The bounds (47) and (48) are obtained as in corollary 3. $\square$

3.1. Numerical experiments

The following tables show numerical experiments on the approximation and the accuracy of the error bounds supplied by corollaries 1-4. In these tables, the second column represents the integral $F_D^r(a, b_1, \dots, b_r; c; x_1, \dots, x_r)$, for $r = 3$ and $r = 4$. The third and sixth columns represent the approximation given by corollaries 1, 2, 3 or 4 for $n = 1$ and $n = 2$ respectively. The fourth and seventh columns represent the respective relative errors, and the fifth and last columns are the respective relative error bounds given in those corollaries.

Parameter values:

$$a = 2, \quad b_1 = 2.25, \quad b_2 = 0.05, \quad b_3 = 0.25, \quad c = 3.5, \quad x_2 = 0.05, \quad x_3 = 0.1, \quad \text{Arg}(x_1) = \frac{-3\pi}{4}$$

$ x_1 $	F_D^3	First or. approx.	Relative error	Relative er. bound	Second or. approx.	Relative error	Relative er. bound
20	0.0084304 - 0.0027687i	0.0011291 - 0.0022707i	0.198	0.388	0.00089615 - 0.0027532i	0.019	0.036
50	0.00012195 - 0.0005966i	0.00014874 - 0.00055883i	0.076	0.126	0.000123723 - 0.0005963i	0.0029	0.0048
100	2.7056e-5 - 0.000175i	3.1065e-5 - 0.0001696i	0.037	0.055	2.718e-5 - 0.0001747i	7.15e-4	1.05e-3
200	5.8769e-6 - 4.9249e-5i	6.4469e-6 - 4.856e-5i	0.018	0.025	5.8855e-6 - 4.9248e-5i	1.74e-4	2.39e-4
500	7.66297e-7 - 8.86727e-6i	8.0748e-7 - 8.8201e-6i	0.007	0.009	7.66537e-7 - 8.86726e-6i	2.7e-5	3.4e-5
1000	1.62758e-7 - 2.37149e-6i	1.682654e-7 - 2.36535e-6i	0.0035	0.0042	1.62774e-7 - 2.37149e-6i	6.6e-6	8.1e-6

Table 1: Approximation supplied by (31) and error bounds given by (33).

Parameter values:

$a = 1.5$, $b_1 = 2.25$, $b_2 = 0.05$, $b_3 = 0.25$, $b_4 = 0.1$, $c = 3.5$, $x_2 = 0.25$, $x_3 = 0.25$, $x_4 = 0.1$, $\text{Arg}(x_1) = \pi$

$ x_1 $	F_D^4	First or. approx.	Relative error	Relative er. bound	Second or. approx.	Relative error	Relative er. bound
20	0.00761345	0.00460439	0.39	0.5	0.00730585	0.04	0.06
50	0.00224492	0.00190431	0.15	0.18	0.00223068	0.0063	0.0096
100	0.0008509	0.00078796	0.074	0.088	0.00084959	0.0015	0.0023
200	0.0003145	0.00030312	0.036	0.042	0.00031442	0.00038	0.00056
500	8.23946e-5	8.1217e-5	0.014	0.016	8.239e-5	6.e-5	8.8e-5
1000	2.9578e-5	2.9383e-5	0.007	0.008	2.9577e-5	1.49e-5	2.2e-5

Table 2: Approximation supplied by (31) and error bounds given by (33).

Parameter values:

$a = 2$, $b_1 = 3$, $b_2 = 0.5$, $b_3 = 0.3$, $c = 4$, $x_2 = 0.2$, $x_3 = 0.3$, $\text{Arg}(x_1) = \pi$

$ x_1 $	F_D^3	First or. approx.	Relative error	Relative er. bound	Second or. approx.	Relative error	Relative er. bound
20	0.00095376	0.0009806	0.028	0.118	0.000956	0.0024	0.01
50	0.00017512	0.00017612	0.0057	0.029	0.00017515	0.0002	0.001
100	4.6326e-5	4.6403e-5	0.00167	0.0099	4.6327e-5	2.87e-5	1.88e-4
200	1.19702e-5	1.19759e-5	4.78e-4	3.46e-3	1.19703e-5	4.12e-6	3.28e-5
500	1.960135e-6	1.960311e-6	9.e-5	8.65e-4	1.960136e-6	3.09e-7	3.29e-6
1000	4.944542e-7	4.944666e-7	2.5e-5	3.04e-4	4.9445423e-7	4.3e-8	5.8e-7

Table 3: Approximation supplied by (39) and error bounds given by $\text{Min}\{(37),(38)\}$.

Parameter values:

$a = 1$, $b_1 = 3$, $b_2 = 0.5$, $b_3 = 0.7$, $b_4 = 0.25$, $c = 5.25$, $x_2 = 0.3$, $x_3 = 0.1$, $x_4 = 0.3$, $\text{Arg}(x_1) = -\frac{\pi}{2}$

$ x_1 $	F_D^4	First or. approx.	Relative error	Relative er. bound	Second or. approx.	Relative error	Relative er. bound
20	0.00320879 - 0.02424707i	0.003088756 - 0.0243536i	0.006	0.037	0.0031949 - 0.0242319i	0.00084	0.0054
50	0.00055794 - 0.0099322i	0.00055321 - 0.009935i	0.0005	0.004	0.0005578 - 0.009932i	2.76e-5	0.0002
100	0.00014352 - 0.00498964i	0.000143146 - 0.0049898i	8.15e-5	6.88e-4	0.000143512 - 0.00498963i	2.03e-6	1.97e-5
200	3.6401e-5 - 0.00249847i	3.6373e-5 - 0.00249848i	1.18e-5	1.22e-4	3.64e-5 - 0.00249847i	1.46e-7	1.75e-6
500	5.875349e-6 - 9.9988e-4i	5.874485e-6 - 9.99988e-4i	9.e-7	1.2e-5	5.875348e-6 - 9.9988e-4i	4.4e-9	7.1e-8
1000	1.473154e-6 - 4.99983e-5i	1.47309e-6 - 4.99983e-5i	1.27e-7	2.19e-6	1.4731537e-6 - 4.99983e-5i	3.05e-10	6.28e-9

Table 4: Approximation supplied by (39) and error bounds given by $\text{Min}\{(37),(38)\}$.

Parameter values:

$$a = 2, b_1 = 1.25, b_2 = 1.05, b_3 = 0.25, c = 3.5, x_3 = 0.1, \text{Arg}(x_1) = \pi, \text{Arg}(x_2) = \pi$$

$ x_1 , x_2 $	F_D^3	First or. approx.	Relative error	Relative er. bound	Second or. approx.	Relative error	Relative er. bound
20, 40	0.00147675	0.00125527	0.15	0.63	0.00146013	0.011	0.057
50, 80	0.000364323	0.000341343	0.063	0.178	0.000363878	0.002	0.0066
200, 100	7.664583e-5	7.460133e-5	0.0267	0.109	7.661635e-5	3.84e-4	0.002
400, 300	1.506288e-5	1.4906e-5	0.01	0.02	1.506206e-5	5.82e-5	1.32e-4
800, 600	4.02922e-6	4.008577e-6	0.005	0.009	4.029166e-6	1.43e-5	3.11e-5
1500, 1000	1.331099e-6	1.327269e-6	2.877e-3	6.45e-3	1.331093e-6	4.6e-6	1.22e-5

Table 5: Approximation supplied by (42) and error bounds given by (44).

Parameter values:

$$a = 1, b_1 = 2 + 0.5i, b_2 = 2 - 1.15i, b_3 = 0.25, b_4 = 0.25, c = 3, x_3 = 0.25, x_4 = 0.1, \text{Arg}(x_1) = \pi, \text{Arg}(x_2) = \pi$$

$ x_1 , x_2 $	F_D^4	First or. approx.	Relative error	Relative er. bound	Second or. approx.	Relative error	Relative er. bound
20, 50	0.00973631 - 0.000433265i	0.00973647 - 0.00043335i	1.9e-5	4.5e-5	0.00973632 - 0.00043327i	1.3e-6	4.8e-6
30, 70	0.006805412 - 0.00033956i	0.00680545 - 0.000339585i	6.5e-6	1.2e-5	0.006805414 - 0.00033956i	3.02e-7	8.1e-7
50, 90	0.00478282 - 0.000331843i	0.00478283 - 0.000331849i	2.06e-6	4.66e-6	0.004782825 - 0.000331843i	6.29e-8	1.99e-7
100, 200	0.002256123 - 1.397424e-4i	0.002256124 - 1.397427e-4i	2.27e-7	2.75e-7	0.002256123 - 1.397424e-4i	3.38e-9	4.3e-9
500, 700	5.5346533e-4 - 4.956593e-5i	5.5346533e-4 - 4.9565931e-5i	3.04e-9	3.87e-9	5.5346533e-4 - 4.956593e-5i	1.1e-11	2.05e-11
1000, 1100	3.1368459e-4 - 3.4142407e-5i	3.1368459e-4 - 3.4142407e-5i	5.4e-10	8.9e-10	3.1368459e-4 - 3.4142407e-5i	2.27e-12	3.e-12

Table 6: Approximation supplied by (42) and error bounds given by $\text{Min}\{(47), (48)\}$.

Parameter values:

$$a = 1, b_1 = 2, b_2 = 1, b_3 = 0.1, c = 2.5, x_3 = 0.25, \text{Arg}(x_1) = \frac{-\pi}{2}, \text{Arg}(x_2) = \pi$$

$ x_1 $	F_D^3	First or. approx.	Relative error	Relative er. bound	Second or. approx.	Relative error	Relative er. bound
30, 40	0.00953984 - 0.0143915i	0.0095395 - 0.0143911i	2.83e-5	0.0015	0.00953984 - 0.0143915i	8.4e-7	0.0001
50, 70	0.0056695 - 0.0084865i	0.0056695 - 0.00848646i	6.69e-6	0.0004	0.0056695 - 0.0084865i	1.22e-7	1.6e-5
100, 150	0.00280578 - 0.00411922i	0.00280578 - 0.0041192i	9.24e-7	5.45e-5	0.00280578 - 0.00411922i	9.34e-9	1.1e-6
400, 300	7.073525e-4 - 0.00140188i	7.073525e-4 - 0.00140188i	3.27e-8	1.5e-5	7.073525e-4 - 0.00140188i	1.9e-10	1.5e-7
700, 600	4.068369e-4 - 7.60952e-4i	4.068369e-4 - 7.60952e-4i	5.86e-9	2.45e-6	4.068369e-4 - 7.60952e-4i	5.38e-11	1.21e-8
1200, 1000	2.369669e-4 - 4.4902823e-5i	2.369667e-4 - 4.4902823e-5i	1.28e-9	6.99e-7	2.369667e-4 - 4.4902823e-5i	1.88e-11	2.08e-9

Table 7: Approximation supplied by (49) and error bounds given by $\text{Min}\{(47), (48)\}$.

Parameter values:

$a = 1$, $b_1 = 2$, $b_2 = 1$, $b_3 = 0.25$, $b_4 = 0.25$, $c = 3.5$, $x_3 = 0.25$, $x_4 = 0.1$, $\text{Arg}(x_1) = \pi$, $\text{Arg}(x_2) = \pi$

$ x_1 $	F_D^4	First or. approx.	Relative error	Relative er. bound	Second or. approx.	Relative error	Relative er. bound
20, 50	0.01672667	0.016724	1.6e-4	0.001	0.0167265	7.44e-6	8.26e-5
30, 70	0.01166452	0.01166385	5.75e-5	4.13e-4	0.0116645	1.65e-6	2.1e-5
50, 90	0.00788302	0.00788289	1.76e-5	2.24e-4	0.00788302	2.6e-7	7.5e-6
100, 200	0.00382053	0.00382052	2.45e-6	2.8e-5	0.00382053	7.86e-9	4.3e-7
500, 700	8.8603367e-4	8.8603364e-4	3.6e-8	1.5e-6	8.8603367e-4	1.6e-10	6.2e-9
1000, 1100	4.834595e-5	4.834596e-5	6.26e-9	5.6e-7	4.834595e-5	3.23e-11	1.5e-9

Table 8: Approximation supplied by (49) and error bounds given by $\text{Min}\{(47),(48)\}$.

4. Acknowledgments

This work has been financially supported by the *DGCYT (BFM2000-0803)* and the *Gobierno de Navarra (Res. 92/2002)* is acknowledged. Special thanks should also be given to the department of mathematics of the University Carlos III of Madrid for their help and hospitality.

References

- [1] A. Al-Zamel, V. K. Tuan, S. L. Kalla, Generalized elliptic-type integrals and asymptotic formulas, *Appl. Math. Comput.* **114**, n.1 (2000) 13-25.
- [2] B.C. CARLSON, The hypergeometric function and the R-function near their branch points, *Rend. Sem. Math. Univ. Politec. Torino*, **Fascicolo speciale** (1985) 63-89.
- [3] B. C. Carlson, Elliptic integrals: symmetric and symbolic integration, *Atti Convegni Lincei*, 147. Rome (1998) 161-181.
- [4] J. Carvalho e Silva and H. M. Srivastava, Asymptotic representations for a class of multiple hypergeometric functions with a dominant variable, *Integral Transform. Spec. Funct.* **11** (2001) 137-150.
- [5] C. Ferreira and J. L. López, Asymptotic expansions of generalized Stieltjes transforms of algebraically decaying functions, *Stud. Appl. Math.* **108** (2002) 187-215
- [6] C.M. Joshi and N.L. Joshi, Expansions of Appell and Lauricella functions, *Tamkang J. of Math.* **27**, n. 4 (1996) 271-279.
- [7] G. Lauricella, Sulle funzioni ipergeometriche a più variabili, *Rend. Circ. mat. Palermo* **7** (1893) 111-158.
- [8] J. L. López, Asymptotic expansions of symmetric standard elliptic integrals, *SIAM J. Math. Anal.* **31**, n. 4 (2000) 754-775.
- [9] J. L. López, Uniform asymptotic expansions of symmetric elliptic integrals, *Const. Approx.* **17**, n. 4 (2001) 535-559.
- [10] J. L. López and C. Ferreira, Asymptotic Expansions of the Generalized Epstein-Hubbell integral, *IMA J. Appl. Math.* **67** (2002) 301-319.
- [11] A. M. Mathai and R. K. Saxena, Various practical problems in probability and statistics where Lauricella's confluent hypergeometric function appears naturally, *Gadnita Sandesh* **1**, n. 1-2 (1987) 41-48.
- [12] Mathur, B. L., On some results involving Lauricella functions, *Bull. Calcutta Math. Soc.* **70**, n. 4 (1978) 221-227.

- [13] W. Miller Jr., Lie theory and Lauricella functions F_D , J. Math Physics **13**, (1972) 1393-1399.
- [14] N. P. Molnar, Approximation of the Lauricella hypergeometric functions F_D^N by branched continued fractions, J. Math. Sci. **90**, n.5 (1998) 2381-2384.
- [15] S. H. Ong and P. A. Lee, Probabilistic interpretations of a transformation of Lauricella's hypergeometric function of n variables and an integral of product of Laguerre polynomials, Int. J. Math. Stat. Sci. **9**, n.1 (2000) 5-13.
- [16] A.P. Prudnikov, Yu.A. Brychkov, O.I. Marichev, Integrals and series, Vol. 1, Gordon and Breach Science Pub., 1990.
- [17] M. Saigo, On properties of the Lauricella hypergeometric function F_D , Bull. Central Res. Inst. Fukuoka Univ. **104**, (1988) 13-31.
- [18] C. K. Sharma and I. J. Singh, Some integrals involving the Lauricella functions and the multivariable H -function, Indian. J. Pure Appl. Math **21**, n.6 (1990) 597-604.
- [19] H.M. Srivastava and P. W. Karlsson, Multiple Gaussian Hypergeometric Series, Wiley, New York, 1985.
- [20] H.M. Srivastava and S.P. Goyal, Fractional Derivatives of the H-function of several variables, J. Math. Anal. Appl., **112**, (1985) 641-651.
- [21] H.M. Srivastava, Some further reduction formulas for certain classes of generalized multiple hypergeometric series arising in physical, astrophysical, and quantum chemical applications, Astrophys. Space Sci., **181**, n.2 (1991) 195-202.
- [22] H.M. Srivastava, Some Lauricella multiple hypergeometric series associated with the product of several Bessel functions, Constantin Carathéodory: an international tribute, Vol I, II, 1304-1341, World Sci. Pub. Teaneck, NJ, 1991.
- [23] N.M. TEMME, Special functions: An introduction to the classical functions of mathematical physics, Wiley and Sons, New York, 1996.
- [24] R. Wong, Distributional derivation of an asymptotic expansion, Proc. Amer. Math. Soc., **80** (1980) 266-270.
- [25] R. Wong, Asymptotic Approximations of Integrals, Academic Press, New York, 1989.