

Elliptic problems involving a gradient term with natural growth

Jean Pierre Gossez

SUMMARY

We consider the problem

$$-\Delta u = g(u)|\nabla u|^2 + f(x, u) \text{ in } \Omega, u > 0. \text{ in } \Omega, u = 0. \text{ on } \partial\Omega,$$

where Ω is a smooth bounded domain in \mathbb{R}^N . Using a Kazdan-Kramer change of variable, this problem can be reduced to a semilinear problem without gradient term, which can then be approachable by variational methods. However various new difficulties arise, in particular with respect to the Ambrosetti-Rabinowitz condition. We investigate some of these difficulties.

In the second part of the talk, we consider the situation of a system. Using again a Kazdan-Kramer change of variables, this system can be transformed into a system without gradient term. In some cases, this latter system can be handled by an upper-lower solution approach, by variational methods, or by blow-up. The blow-up procedure leads to the study of new Liouville type theorems.

Joint work with D. de Figueiredo (Campinas, Brasil), H. Ramos Quoirin (Cordoba, Argentina) and P. Ubilla (Santiago, Chile).

¹Département de mathématique, CP214, Campus de la Plaine,
Université Libre de Bruxelles, 1050 Bruxelles, Belgique
email: gossez@ulb.ac.be